

Director of Public Health Annual Report 2005: Choosing Health in Leicester

Health Facts 4 - Cancer rates

Source : National Centre for Health Outcomes Development, NHS Health and Social Care Information Centre

Males

Cancer	ICD 10	New cases in Leicester (1998-2000)	Deaths in Leicester 2003	SRR	Standardised mortality ratio (all ages, 2001-2003)				Rate of FYLL per 100,000 population (age standardised, 2001-2003)					1 yr survival LNR	5 yr survival LNR
					East Mids	Leicester	EL PCT	LCW PCT	England	East Mids	Leicester	EL PCT	LCW PCT		
All	C00-C97		318		97*	101	93	113**	171.2	163.9	171.8	154.4	199.7		
Lung	C33-C34	305	76	105	95*	104	88	130**	38.7	35.6	45.1	35.6	59.8	19%	4.1%
Colorectal	C17-C21	213	26	90	100	102	95	113	17.4	17.4	16.8	14	21.7	63.7%	42.9%
Stomach	C16	107	16	138**	104	126	107	154**	6.7	6.9	8.5	4.2	15.6	30.6%	8.5%
Oesophageal	C15	49	13	92	104	98	88	114	11.4	12.6	7.7	5.2	11.8	22.6%	4.9%
Bladder	C67	93	17	86	101	98	80	125	3.8	3.5	2.1	0.9	3.9	83%	68.3%
Malignant melanoma	C43	22	5	62*	87	93	77	118	4	3.8	1.4	0.7	2.4		
Prostate	C61	236	40	73*	100	92	89	98	7.4	7.2	8.5	11.9	3.3	83%	56.9%
Leukaemia	C91-C95		18		101	102	107	94	8.1	8.2	9.3	9.2	9.9		

SRR = standardised registration ratio. Number of cancers registered in a population compared with national rates

Standardised mortality ratio - see glossary

FYLL = future years of life lost - see glossary

** = significantly above national rates

* = significantly below national rates

Cancer rates Health Facts from annual reports 2005 to 2010

Females

Cancer	ICD 10	New cases in Leicester (1998-2000)	Deaths in Leicester 2003	SRR	Standardised mortality ratio (all ages, 2001-2003)				Rate of FYLL per 100,000 population (age standardised, 2001-2003)					1 yr survival LNR	5 yr survival LNR
					East Mids	Leicester	EL PCT	LCW PCT	England	East Mids	Leicester	EL PCT	LCW PCT		
All	C00-C97		289		100	97	86*	111**	157.7	158.4	157.2	146.2	174.3		
Lung	C33-C34	189	58	106	95*	105	77*	145**	23.5	22.5	24.4	16.9	36	20.7%	5.7%
Colorectal	C17-C21	184	28	87	100	82	66*	103	10.9	11.7	8.9	6.9	12.1	62.6%	41.7%
Stomach	C16	45	14 (2002)	104	102	114	130	93	2.9	2.8	4.9	4.4	5.5	28.6%	10.5%
Oesophageal	C15	46	10	131	106	101	102	101	3.3	3.4	2.3	2.6	1.9	25.6%	4.8%
Bladder	C67	50	6	115	104	88	77	104	1.5	1.6	1.7	1.9	1.2	74.1%	59.8%
Malignant melanoma	C43	25	4	53*	85	61	70	50	2.9	2.5	0.9	0.2	2		
Breast	C50	457	54	91*	102	105	91	125	39.6	40.2	42.7	38.6	49.3	91.3%	73.7%
Leukaemia	C91-C95		7		99	109	93	130	5.5	4.8	9.0	10.3	6.9		
Cervical	C53	41	9	100	99	103	99	108	5.2	5.2	5.8	5.9	5.5	78.9%	59.7%

SRR = standardised registration ratio. Number of cancers registered in a population compared with national rates

Standardised mortality ratio - see glossary

FYLL = future years of life lost - see glossary

** = significantly above national rates

* = significantly below national rates

Choosing Health in Leicester's Localities, The Director of Public Health Annual Report 2006

Health Facts 4 - Cancer rates

Source: National Centre for Health Outcomes Development, NHS Health and Social Care Information Centre

Cancer rate in Males

Cancer	ICD 10	New cases In Leicester (2000-02)	SRR Leicester (2000-02)	No. deaths in Leicester (2004)	Standardised Mortality Ratio (Indirect) for all ages, 2002-04				Rate of YLL per 10,000 population(age standardised) 2002-2004					1 yr survival LNR	5 yr survival LNR
			Leicester SMR		E. Mids SMR	Leicester SMR	ELPCT SMR	LCWPCT SMR	England SYLL	E. Mids SYLL	Leicester SYLL	ELPCT SYLL	LCWPCT SYLL		
All	C00-C97	-	-	286	98.2	97.9	90.7	108.8	167.0	161.7	166.8	146.8	199.8	-	-
Lung	C33-C34	301	112.0	62	95.5	94.0	83.0	110.8	37.3	34.5	36.4	32.8	42.4	20.2%	4.7%
Colorectal	C17-C21	207	91.0	31	102.1	98.6	90.9	110.3	16.8	16.9	15.1	13.3	18.6	64.9%	42.6%
Stomach	C16	107	152.2	22	103.4	146.2	146.5	145.8	6.3	6.7	10.8	7.8	15.8	29.8%	8.6%
Oesophageal	C15	57	104.5	10	100.7	84.7	56.0	128.6	11.4	12.0	6.9	2.4	14.0	24.9%	5.5%
Bladder	C67	70	77.7	9	100.5	95.4	90.4	102.8	3.7	3.3	1.8	1.4	2.5	83.1%	72.8%
Malignant Melanoma	C43	32	78.4	1	86.1	63.0	74.0	45.9	4.1	4.0	0.5	0.7	0.0		
Prostate	C61	308	82.7	41	101.6	89.9	94.5	83.0	7.3	7.5	9.2	12.1	4.7	84.8%	60.2%
Leukaemia	C91-C95	-	-	9	102.5	117.2	108.6	130.3	7.6	7.6	9.3	8.2	11.4	-	-
Hodgkins	C81	-	-	0	96.7	0.0	0.0	0.0	1.0	0.7	0.0	0.0	0.0	-	-

Significance of Years of Life Lost not shown

- significantly worse than the national rate (100)
- significantly better than the national rate (100)

Cancer rate in Females

Cancer	ICD 10	New cases in Leicester (2000-02)	SRR Leicester (2000-02) Leicester SMR	No. deaths in Leicester (2004)	Standardised Mortality Ratio (Indirect) for all ages, 2002-04				Rate of YLL per 10,000 population(age standardised) 2002-2004					1 yr survival	5 yr survival
					E. Mids SMR	Leicester SMR	ELPCT SMR	LCWPCT SMR	England SYLL	E. Mids SYLL	Leicester SYLL	ELPCT SYLL	LCWPCT SYLL	LNR	LNR
All	C00-C97	-	-	299	99.5	99.6	89.1	115.0	154.3	152.9	154.3	129.8	191.4	-	-
Lung	C33-C34	205	116.0	50	93.8	107.1	80.7	144.5	23.6	21.7	24.6	15.9	38.3	20.6%	6.2%
Colorectal	C17-C21	191	96.0	30	103.0	86.3	85.9	86.9	10.7	11.4	6.3	5.3	7.7	63.6%	42.4%
Stomach	C16	49	123.7	9	100.0	104.9	123.1	79.9	2.9	2.7	3.7	4.3	2.7	29.4%	14.0%
Oesophageal	C15	38	113.4	9	105.2	99.1	91.2	110.0	3.3	3.5	1.6	1.5	1.8	24.8%	6.8%
Bladder	C67	30	82.3	5	105.4	74.9	78.1	70.5	1.5	1.6	0.8	0.7	1.0	75.3%	59.5%
Malignant Melanoma	C43	38	70.7	0	88.6	41.5	52.8	25.3	3.0	2.6	0.2	0.2	0.1	-	-
Breast	C50	512	101.9	59	100.2	108.5	88.3	137.5	38.3	38.8	40.1	34.8	48.1	92.0%	76.1%
Cervical	C53	58	152.8	10	97.7	158.7	140.4	185.2	5.0	4.5	9.7	7.2	12.8	-	-
Leukaemia	C91-C95	-	-	2	98.2	84.2	79.5	90.8	5.1	3.8	6.9	8.7	4.1	80.6%	58.3%
Hodgkins	C81	-	-	1	105.7	180.1	101.5	294.2	0.7	1.0	1.1	0.1	2.6	-	-

Significance of Years of Life Lost not shown

- significantly worse than the national rate (100)
- significantly better than the national rate (100)

ICD 10: International Classification of Diseases: WHO's internationally accepted classification of death and disease, revision 10.

Standardised Registration Ratio (SRR) (Cancer): Ratio of cancers registered in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability

Standardised Mortality Ratio: Ratio of number of deaths in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability.

Years of life lost rate: Potential number of years of life lost as a result of premature death (under 75 years). Directly age-standardised per 100,000 standard European population

Survival rate: Ratio of the survival rate actually observed among the (cancer) patients and the survival that would have been expected if they had only had the same overall mortality rates as the general population Survival rate - 1 year / 5 years: Relative survival rate observed at one and five years after diagnosis, compared with general population.

Health in Leicester Annual Report of the Director of Public Health and Health Improvement 2007

Health Facts 4 - Cancer rates

Source: National Centre for Health Outcomes Development, NHS Health and Social Care Information Centre

Cancer rate in Males

Cancer	ICD 10	New cases in Leicester (2002-04)	SRR Leicester (2002-04)	No. deaths in Leicester (2006)	Standardised Mortality Ratio (Indirect), 2004-06 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2004-06), all ages			Directly age-standardised mortality rate per 100,000 (2004-06), under 75s			1 yr survival (1997-99)	5 yr survival (1997-99)
			Leicester SRR		England SMR	E.Mids SMR	Leicester SMR	England DSR	E.Mids DSR	Leicester DSR	England DSR	E.Mids DSR	Leicester DSR	LNR	LNR
All	C00-C97	1537	94.0	282	100	98.4	96.3	213.7	209.5	206.5	130.4	126.6	128.4	-	-
Lung	C33-C34	270	107.3	74	100	95.4	100.7	51.1	48.7	51.6	33.1	31.0	32.6	20.6%	5.3%
Colorectal	C17-C21	194	86.3	27	100	97.5	79.8	23.0	22.3	18.2	13.9	13.2	10.9	64.6%	44.9%
Stomach	C16	81	129.9	14	100	100.0	138.2	8.9	8.9	12.7	5.1	5.5	9.6	33.7%	9.6%
Oesophageal	C15	53	95.8	16	100	99.0	76.7	13.0	12.8	10.3	9.0	8.6	7.4	28.5%	7.0%
Bladder	C67	77	92.2	14	100	99.6	101.3	8.2	8.1	8.5	3.7	3.4	4.1	83.4%	71.6%
Malignant Melanoma	C43	21	46.4	4	100	93.3	67.8	2.9	2.7	2.0	2.3	2.1	1.2	-	-
Prostate	C61	306	78.5	21	100	100.7	87.3	25.7	25.9	22.2	9.1	9.2	7.5	85.2%	63.6%
Leukaemia	C91-C95	-	-	6	100	100.4	101.9	6.7	6.6	6.6	4.2	4.3	4.5	-	-
Hodgkins	C81	-	-	0	100	91.4	45.3	0.5	0.5	0.2	0.4	0.4	0.2	-	-

Significance of Years of Life Lost not shown

significantly worse than the national rate (100)
 significantly better than the national rate (100)

Cancer rate in Females

Cancer	ICD 10	New cases in Leicester (2002-04)	SRR Leicester (2002-04)	No. deaths in Leicester (2006)	Standardised Mortality Ratio (Indirect), 2004-06 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2004-06), all ages			Directly age-standardised mortality rate per 100,000 (2004-06), under 75s			1 yr survival (1997-99)	5 yr survival (1997-99)
			Leicester SRR		England SMR	E.Mids SMR	Leicester SMR	England SR	E.Mids DSR	Leicester DSR	England DSR	E.Mids DSR	Leicester DSR	LNR	LNR
All	C00-C97	1674	100.8	279	100	99.0	103.2	152.2	150.9	157.1	105.1	104.5	107.4	-	-
Lung	C33-C34	207	118.3	48	100	89.9	94.2	28.9	26.0	27.3	20.5	18.7	19.4	23.3%	8.2%
Colorectal	C17-C21	188	96.8	29	100	97.6	94.0	14.4	14.3	13.3	8.6	8.8	7.3	63.6%	41.0%
Stomach	C16	41	114.7	2	100	94.0	89.3	3.8	3.6	3.9	2.1	2.1	2.9	30.8%	12.8%
Oesophageal	C15	33	103.6	18	100	107.2	129.8	4.8	5.1	6.4	2.8	3.0	4.0	28.2%	*5.7%
Bladder	C67	27	79.1	5	100	104.9	112.1	2.8	2.9	2.9	1.4	1.4	1.0	79.0%	66.2%
Malignant Melanoma	C43	33	56.5	1	100	95.7	42.1	1.9	1.9	0.8	1.5	1.6	0.8	-	-
Breast	C50	500	96.2	51	100	102.9	111.3	28.0	28.7	30.7	21.6	21.9	22.6	92.5%	77.1%
Cervical	C53	42	115.6	5	100	87.7	158.5	2.5	2.1	4.4	2.1	1.7	3.9	84.7%	61.6%
Leukaemia	C91-C95	-	-	5	100	100.1	49.5	3.8	3.7	2.0	2.4	2.3	1.2	-	-
Hodgkins	C81	-	-	1	100	108.3	120.4	0.3	0.3	0.3	0.3	0.2	0.0	-	-

Significance of Years of Life Lost not shown

■ significantly worse than the national rate (100)

■ significantly better than the national rate (100)

ICD 10: International Classification of Diseases: WHO's internationally accepted classification of death and disease, revision 10.

Standardised Registration Ratio (SRR) (Cancer): Ratio of cancers registered in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability

Standardised Mortality Ratio: Ratio of number of deaths in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability.

Years of life lost rate: Potential number of years of life lost as a result of premature death (under 75 years). Directly age-standardised per 100,000 standard European population

Survival rate: Ratio of the survival rate actually observed among the (cancer) patients and the survival that would have been expected if they had only had the same overall mortality rates as the general population

Survival rate - 1 year / 5 years: Relative survival rate observed at one and five years after diagnosis, compared with general population

Improving Health in Leicester: Annual Report of the Director of Public Health and Health Improvement 2008/09

Health Facts 4: Cancer rates

Source: National Centre for Health Outcomes Development, NHS Health and Social Care Information Centre

Cancer rates in males

Cancer	ICD 10	New cases in Leicester (2002-04)	SRR Leicester (2002-04)	No. deaths in Leicester (2006)	Standardised Mortality Ratio (Indirect), 2004-6 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2004-6), All ages			Directly age-standardised mortality rate per 100,000 (2004-6), under 75s			1 yr survival (1997-99)	5 yr survival (1997-99)
			Leicester SRR		England SMR	E.Mids SMR	Leicester SMR	England DSR	E.Mids DSR	Leicester DSR	England DSR	E.Mids DSR	Leicester DSR	LNR	LNR
All	C00-C97	1537	94.0	282	100	98.4	96.3	213.7	209.5	206.5	130.4	126.6	128.4	-	-
Lung	C33-C34	270	107.3	74	100	95.4	100.7	51.1	48.7	51.6	33.1	31.0	32.6	20.6%	5.3%
Colorectal	C17-C21	194	86.3	27	100	97.5	79.8	23.0	22.3	18.2	13.9	13.2	10.9	64.6%	44.9%
Stomach	C16	81	129.9	14	100	100.0	138.2	8.9	8.9	12.7	5.1	5.5	9.6	33.7%	9.6%
Oesophageal	C15	53	95.8	16	100	99.0	76.7	13.0	12.8	10.3	9.0	8.6	7.4	28.5%	7.0%
Bladder	C67	77	92.2	14	100	99.6	101.3	8.2	8.1	8.5	3.7	3.4	4.1	83.4%	71.6%
Malignant Melanoma	C43	21	46.4	4	100	93.3	67.8	2.9	2.7	2.0	2.3	2.1	1.2	-	-
Prostate	C61	306	78.5	21	100	100.7	87.3	25.7	25.9	22.2	9.1	9.2	7.5	85.2%	63.6%
Leukaemia	C91-C95	-	-	6	100	100.4	101.9	6.7	6.6	6.6	4.2	4.3	4.5	-	-
Hodgkins	C81	-	-	0	100	91.4	45.3	0.5	0.5	0.2	0.4	0.4	0.2	-	-

- Significantly better than the national rate (100)
- Significantly worse than the national rate (100)

Cancer rates in females

Cancer	ICD 10	New cases in Leicester (2002-04)	SRR Leicester (2002-04)	No. deaths in Leicester (2006)	Standardised Mortality Ratio (Indirect), 2004-6 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2004-6), All ages			Directly age-standardised mortality rate per 100,000 (2004-6), under 75s			1 yr survival (1997-99)	5 yr survival (1997-99)
			Leic SRR		England SMR	E.Mid SMR	Leicester SMR	England DSR	E. Mids DSR	Leicester DSR	England DSR	E. Mids DSR	Leicester DSR	LNR	LNR
All	C00-C97	1674	100.8	279	100	99.0	103.2	152.2	150.9	157.1	105.1	104.5	107.4	-	-
Lung	C33-C34	207	118.3	48	100	89.9	94.2	28.9	26.0	27.3	20.5	18.7	19.4	23.3%	8.2%
Colorectal	C17-C21	188	96.8	29	100	97.6	94.0	14.4	14.3	13.3	8.6	8.8	7.3	63.6%	41.0%
Stomach	C16	41	114.7	2	100	94.0	89.3	3.8	3.6	3.9	2.1	2.1	2.9	30.8%	12.8%
Oesophageal	C15	33	103.6	18	100	107.2	129.8	4.8	5.1	6.4	2.8	3.0	4.0	28.2%	*5.7%
Bladder	C67	27	79.1	5	100	104.9	112.1	2.8	2.9	2.9	1.4	1.4	1.0	79.0%	66.2%
Malignant Melanoma	C43	33	56.5	1	100	95.7	42.1	1.9	1.9	0.8	1.5	1.6	0.8	-	-
Breast	C50	500	96.2	51	100	102.9	111.3	28.0	28.7	30.7	21.6	21.9	22.6	92.5%	77.1%
Cervical	C53	42	115.6	5	100	87.7	158.5	2.5	2.1	4.4	2.1	1.7	3.9	84.7%	61.6%
Leukaemia	C91-C95	-	-	5	100	100.1	49.5	3.8	3.7	2.0	2.4	2.3	1.2	-	-
Hodgkins	C81	-	-	1	100	108.3	120.4	0.3	0.3	0.3	0.3	0.2	0.0	-	-

*not age- standardised

- Significantly better than the national rate (100)
- Significantly worse than the national rate (100)

ICD 10: International Classification of Diseases: WHO's internationally accepted classification of death and disease, revision 10.

SRR Standardised Registration Ratio: Ratio of cancers registered in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability.

SMR Standardised Mortality Ratio: Ratio of number of deaths in a population compared with the national population, standardised to adjust for differences in age and sex of the local population. A score greater than 100 indicates an increased probability and a score below 100 indicates a reduced probability.

Standardised Years of Life Lost Rate: Potential number of years of life lost as a result of premature death (under 75 years) per 10,000 European standard population

Survival Rate: Ratio of the survival rate actually observed among the cancer patients and the survival that would have been expected if they had only had the same overall mortality rates as the general population

Survival Rate: 1 year/ 5years: Relative survival rate observed at one and five years after diagnosis, compared with general population



Improving Health in Leicester: Annual Report of the Director of Public Health and Health Improvement 2010

Health Facts 4: Cancer rates

Source: National Centre for Health Outcomes Development, NHS Health and Social Care Information Centre

Cancer rates in males

Cause of death	ICD 10	New cases in Leicester (2004-06)	No. deaths in Leicester (2008)	Standardised Mortality Ratio (Indirect), 2006-8 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2006-8), All ages			Directly age-standardised mortality rate per 100,000 (2004-6), under 75s					1 yr survival (2000-02)	5 yr survival (2000-02)
				England	East Mids	Leicester	England	East Mids	Leicester	England	East Mids	Leicester				
				SMR	SMR	SMR	DSR	DSR	DSR	DSR	DSR	DSR	LCL	UCL		
All	C00-C97	1639	303	100	99.9	100.3	207.4	207.1	206.2	126.3	125.2	122.2	110.6 2	133.75	-	-
Lung	C33-C34	320	82	100	97.1	110.0	49.7	48.1	54.7	31.9	30.7	34.5	28.27	40.68	25.4%	6.7%
Colorectal	C17- C21	211	36	100	98.7	91.0	22.2	21.9	20.4	13.4	13.1	12.8	9.05	16.56	66.9%	47.9%
Stomach	C16	64	19	100	102.4	139.1	8.0	8.2	11.4	4.5	4.6	7.5	4.60	10.36	34.1%	12.3%
Oesophageal	C15	66	15	100	104.5	99.7	12.8	13.4	12.1	8.9	9.3	6.9	4.10	9.60	35.2%	10.5%
Bladder	C67	72	10	100	102.4	95.8	8.0	8.1	7.5	3.6	3.5	3.1	1.25	4.89	79.5%	62.1%
Malignant Melanoma	C43	29	1	100	92.0	55.2	2.9	2.7	2.0	2.5	2.4	1.1	0.01	2.20	-	-
Prostate	C61	332	25	100	104.5	68.7	24.5	25.6	16.6	8.7	8.9	4.7	2.37	6.95	88.7%	69.4%
Leukaemia	C91- C95	-	10	100	100.9	70.2	6.6	6.7	4.4	4.1	4.2	2.4	0.83	3.99	-	-
Hodgkins	C81	-	2	100	102.1	144.9	0.5	0.5	0.7	0.4	0.4	0.5	0.00	1.25		

 Significantly better than the national rate (100)
 Significantly worse than the national rate (100)

Cancer rates in females

Cause of death	ICD 10	New cases in Leicester (2004-06)	No. deaths in Leicester (2008)	Standardised Mortality Ratio (Indirect), 2006-8 pooled, all ages			Directly age-standardised mortality rate per 100,000 (2006-8), All ages			Directly age-standardised mortality rate per 100,000 (2004-6), under 75s					1 yr survival (2000-02) EM SHA	5 yr survival (2000-02) EM SHA
				England	East Mids	Leicester	England	East Mids	Leicester	England	East Mids	Leicester				
				SMR	SMR	SMR	DSR	DSR	DSR	DSR	DSR	DSR	LCL	UCL		
All	C00-C97	1713	297	100	99.3	105.5	149.9	149.2	159.5	102.7	102.4	108.3	97.61	118.92	-	-
Lung	C33-C34	199	54	100	93.1	113.2	30.0	28.3	34.3	21.1	20.1	24.5	19.41	29.54	26.6%	7.8%
Colorectal	C17- C21	176	29	100	97.6	105.3	14.1	13.7	14.8	8.4	8.1	8.7	5.73	11.73	67.1%	48.6%
Stomach	C16	23	13	100	101.0	94.0	3.5	3.5	3.5	1.9	2.0	2.3	0.80	3.85	31.3%	15.5%
Oesophageal	C15	39	13	100	110.4	153.9	4.7	5.1	7.5	2.8	3.0	4.9	2.65	7.05	32.4%	10.4%
Bladder	C67	33	3	100	105.4	87.0	2.8	2.9	1.9	1.3	1.3	0.2	0.00	0.67	68.2%	47.8%
Malignant Melanoma	C43	32	4	100	99.0	49.4	2.0	2.1	0.6	1.6	1.7	0.2	0.00	0.67	-	-
Breast	C50	543	45	100	103.3	100.7	26.8	27.4	28.6	20.6	20.8	22.8	17.90	27.78	93.9%	79.7%
Cervical	C53	48	4	100	95.7	156.9	2.3	2.2	3.8	2.0	1.9	3.2	1.39	5.09	80.6%	65.2%
Leukaemia	C91- C95	-	8	100	93.3	93.0	3.7	3.5	3.2	2.3	2.2	1.6	0.31	2.91	-	-
Hodgkins	C81	-	2	100	93.5	219.3	0.4	0.3	0.6	0.3	0.3	0.3	0.00	0.75	-	-

*not age-standardised

- Significantly better than the national rate (100)
- Significantly worse than the national rate (100)